

# Azygos Vein, Hemiazygos Vein, Accessory Hemiazygos Vein, Thoracic Sympathetic Trunk

---

## Azygos Vein

- The **azygos vein** drains the **thoracic wall** and **upper lumbar region** and provides an important connection between the **superior and inferior vena cava**, as well as the **portal and vertebral venous systems**.
- **Formation:**
  - Formed by the union of **right subcostal vein** and **right ascending lumbar vein**.
  - Sometimes includes a contribution from the **lumbar azygos vein**.
- **Course:**
  - Enters the thorax through the **aortic opening** of the diaphragm.
  - Ascends on the **right side of the vertebral column** and arches over the **root of the right lung** to join the **superior vena cava**.
- **Relations:**
  - **Anteriorly:** Oesophagus.
  - **Posteriorly:** Lower eight thoracic vertebrae and right posterior intercostal arteries.
  - **Right side:** Right lung and pleura, greater splanchnic nerve.

- **Left side:** Thoracic duct and aorta below, oesophagus and vagus nerve above.

- **Tributaries:**

1. Right superior intercostal vein (from 2nd–4th intercostal spaces).
2. 5th–11th right posterior intercostal veins.
3. Hemiazygos and accessory hemiazygos veins.
4. Right bronchial vein.
5. Oesophageal, mediastinal, and pericardial veins.

- **Clinical Note:**

- Acts as an important **collateral channel** during **obstruction of the inferior vena cava**

bd-chaurasias-human-anatomy-vol...

---

## Hemiazygos Vein

- Also known as the **inferior hemiazygos vein**, it mirrors the **lower part** of the azygos vein on the left side.

- **Formation:**

- Union of **left ascending lumbar vein**, **left subcostal vein**, and **left lumbar azygos vein**.

- **Course:**

- Enters thorax by piercing the **left crus of diaphragm**, ascends along the **left side of vertebral column** behind the **aorta**, and at the level of the **8th thoracic vertebra**, turns to the right behind the **oesophagus** and **thoracic duct** to join the **azygos vein**.

- **Tributaries:**

- 9th–11th left posterior intercostal veins.
- Oesophageal veins.

- **Function:**

- Provides venous drainage for the **lower left posterior thoracic wall** and connects with the **azygos system** for collateral circulation  
bd-chaurasias-human-anatomy-vol...

.

---

## Accessory Hemiazygos Vein

- Also called the **superior hemiazygos vein**, representing the **upper counterpart** of the hemiazygos vein.

- **Formation and Course:**

- Begins around the **4th or 5th intercostal space**, descends along the **left side of vertebral column**, and at the level of the **8th thoracic vertebra**, crosses to the right behind the **aorta** and **thoracic duct** to join the **azygos vein**.

- **Tributaries:**

- 5th–8th left posterior intercostal veins.

- Sometimes **left bronchial veins**.

- **Function:**

- Drains the **upper left posterior thoracic wall** and provides a connection between the **superior intercostal** and **hemiazygos systems**

bd-chaurasias-human-anatomy-vol...

.

---

## Thoracic Sympathetic Trunk

- The **thoracic part of the sympathetic trunk** is a **chain of ganglia** lying on each side of the **vertebral column**.

- **Extent:**

- Continuous above with the **cervical** and below with the **lumbar sympathetic trunks**.

- **Structure:**

- Usually **11 ganglia**, corresponding to **T1–T12** spinal nerves.

- The **first thoracic ganglion** often fuses with the **inferior cervical ganglion** to form the **stellate (cervicothoracic) ganglion**.

- **Course and Relations:**

- Descends across the **neck of the 1st rib**, heads of 2nd–10th ribs, and bodies of 11th–12th thoracic vertebrae.

- Lies anterior to **posterior intercostal vessels** and **nerves**; continues below behind the **medial arcuate ligament** into the abdomen.

- **Branches:**

1. **Grey rami communicantes** – to all thoracic spinal nerves (T1–T12), carrying **postganglionic fibres** to blood vessels, sweat glands, and arrector pili.
2. **White rami communicantes** – from T1–T12 to corresponding ganglia; T1–T5 ascend to cervical ganglia, T10–L2 descend to lumbar or sacral ganglia.
3. **Visceral branches:**

- **Cardiac, pulmonary, and oesophageal branches** (from upper 5 ganglia).
- **Splanchnic nerves** from lower 8 ganglia:
  - **Greater splanchnic nerve (T5–T9)**
  - **Lesser splanchnic nerve (T10–T11)**
  - **Least splanchnic nerve (T12)**

- **Clinical Significance:**

- Involved in **autonomic control of thoracic viscera**.
- **Stellate ganglion block** is performed for **relief of vasospasm or hyperhidrosis**.
- Damage can cause **Horner's syndrome** (ptosis, miosis, anhidrosis)